Product Name: Integrin α1 Rabbit Polyclonal Antibody Catalog #: APRab12665



Summary

Production Name Integrin α1 Rabbit Polyclonal Antibody

Description Rabbit Polyclonal Antibody

Host Rabbit
Application IHC,ELISA

Reactivity Human, Rat, Mouse

Performance

ConjugationUnconjugatedModificationUnmodified

Isotype IgG

Clonality Polyclonal Form Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw

cycles.

Buffer Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.

Purification Affinity purification

Immunogen

Storage

Gene Name ITGA1

Integrin alpha-1 (CD49 antigen-like family member A;Laminin and collagen Alternative Names

receptor; VLA-1; CD antigen CD49a)

Gene ID 3672.0

SwissProt ID P56199.Synthetic peptide from human protein at AA range: 920-980

Application

Dilution Ratio IHC 1:50-200, ELISA 1:10000-20000.

Molecular Weight

Background

Catalog #: APRab12665




integrin subunit alpha 1(ITGA1) Homo sapiens This gene encodes the alpha 1 subunit of integrin receptors. This protein heterodimerizes with the beta 1 subunit to form a cell-surface receptor for collagen and laminin. The heterodimeric receptor is involved in cell-cell adhesion and may play a role in inflammation and fibrosis. The alpha 1 subunit contains an inserted (I) von Willebrand factor type I domain which is thought to be involved in collagen binding. [provided by RefSeq, Jul 2008], domain: The integrin I-domain (insert) is a VWFA domain. Integrins with I-domains do not undergo protease cleavage, function: Integrin alpha-1/beta-1 is a receptor for laminin and collagen. It recognizes the proline-hydroxylated sequence G-F-P-G-E-R in collagen., similarity: Belongs to the integrin alpha chain family., similarity: Contains 1 VWFA domain., similarity: Contains 7 FG-GAP repeats., subunit: Heterodimer of an alpha and a beta subunit. Alpha-1 associates with beta-1. Interacts with RAB21.,


Research Area

Focal adhesion; ECM-receptor interaction; Hematopoietic cell lineage; Regulates Actin and Cytoskeleton; Hypertrophic cardiomyopathy (HCM); Arrhythmogenic right ventricular cardiomyopathy (ARVC); Dilated cardiomyopathy;

Image Data



Immunohistochemical analysis of paraffin-embed ast-cancer, antibody was diluted at 1:200




Immunohistochemical analysis of paraffin-embedded human-colon-cancer, antibody was diluted at 1:200


Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Catalog #: APRab12665






Immunohistochemical analysis of paraffin-embedded human-colon-cancer, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-lung-cancer, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-lung-cancer, antibody was diluted at 1:200

Note

For research use only.