
Summary

Production Name	Laminin γ -1 Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human,Mouse,Rat,Monkey,Cat

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	LAMC1
Alternative Names	LAMC1; LAMB2; Laminin subunit gamma-1; Laminin B2 chain; Laminin-1 subunit gamma; Laminin-10 subunit gamma; Laminin-11 subunit gamma; Laminin-2 subunit gamma; Laminin-3 subunit gamma; Laminin-4 subunit gamma; Laminin-6 subunit gamma; Lamini
Gene ID	3915.0
SwissProt ID	P11047.The antiserum was produced against synthesized peptide derived from human LAMC1. AA range:1451-1500

Application

Dilution Ratio	WB 1:500 - 1:2000. ELISA: 1:40000
Molecular Weight	178kD


Background

Laminins, a family of extracellular matrix glycoproteins, are the major noncollagenous constituent of basement membranes. They have been implicated in a wide variety of biological processes including cell adhesion, differentiation, migration, signaling, neurite outgrowth and metastasis. Laminins, composed of 3 non identical chains: laminin alpha, beta and gamma (formerly A, B1, and B2, respectively), have a cruciform structure consisting of 3 short arms, each formed by a different chain, and a long arm composed of all 3 chains. Each laminin chain is a multidomain protein encoded by a distinct gene. Several isoforms of each chain have been described. Different alpha, beta and gamma chain isomers combine to give rise to different heterotrimeric laminin isoforms which are designated by Arabic numerals in the order of their discovery, i.e. alpha1beta1gamma1 heterotrimer is laminin 1. The biological function: Domains VI and IV are globular, domain: The alpha-helical domains I and II are thought to interact with other laminin chains to form a coiled coil structure, function: Binding to cells via a high affinity receptor, laminin is thought to mediate the attachment, migration and organization of cells into tissues during embryonic development by interacting with other extracellular matrix components, similarity: Contains 1 laminin IV type A domain, similarity: Contains 1 laminin N-terminal domain, similarity: Contains 11 laminin EGF-like domains, subunit: Laminin is a complex glycoprotein, consisting of three different polypeptide chains (alpha, beta, gamma), which are bound to each other by disulfide bonds into a cross-shaped molecule comprising one long and three short arms with globules at each end. Gamma-1 is a subunit of laminin-1 (EHS laminin), laminin-2 (merosin), laminin-3 (S-laminin), laminin-4 (S-merosin), laminin-6 (K-laminin) and laminin-7 (KS-laminin), tissue specificity: Found in the basement membranes (major component),

Research Area

Focal adhesion; ECM-receptor interaction; Prion diseases; Pathways in cancer; Small cell lung cancer;

Image Data




Western blot analysis of lysates from HUVEC cells, using LAMC1 Antibody. The lane on the right is blocked with the synthesized peptide.

Product Name: Laminin γ -1 Rabbit Polyclonal Antibody
Catalog #: APRab13203



customer's (cat sample)



Western Blot analysis of customer 's (cat sample) using Laminin γ -1 Polyclonal Antibody diluted at 1: 1000

Note

For research use only.