

Summary

Production Name	LPCAT2 Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB
Reactivity	Human,Mouse,Rat

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	LPCAT2 LPCAT2; AYTL1; Lysophosphatidylcholine acyltransferase 2; LPC acyltransferase 2;
Alternative Names	LPCAT-2; LysoPC acyltransferase 2; 1-acylglycerophosphocholine O-acyltransferase; 1-alkylglycerophosphocholine O-acetyltransferase; Acetyl-CoA:lyso-platelet-ac
Gene ID	54947.0
SwissProt ID	Q7L5N7.The antiserum was produced against synthesized peptide derived from human LPCAT2. AA range:321-370

Application


Dilution Ratio	WB 1:500-2000
Molecular Weight	50kD

Background

This gene encodes a member of the lysophospholipid acyltransferase family. The encoded enzyme may function in two ways: to catalyze the biosynthesis of platelet-activating factor (1-O-alkyl-2-acetyl-sn-glycero-3-phosphocholine) from 1-O-alkyl-sn-glycero-3-phosphocholine, and to catalyze the synthesis of glycerophospholipid precursors from arachidonyl-CoA and lysophosphatidylcholine. The encoded protein may function in membrane biogenesis and production of platelet-activating factor in inflammatory cells. The enzyme may localize to the endoplasmic reticulum and the Golgi. [provided by RefSeq, Feb 2009],catalytic activity:Acetyl-CoA + 1-alkyl-sn-glycero-3-phosphocholine = CoA + 2-acetyl-1-alkyl-sn-glycero-3-phosphocholine.,catalytic activity:Acyl-CoA + 1-acyl-sn-glycero-3-phosphocholine = CoA + 1,2-diacyl-sn-glycero-3-phosphocholine.,domain:The HXXXXD motif is essential for acyltransferase activity.,enzyme regulation:Acetyltransferase activity is increased following acute inflammatory stimulation by lipopolysaccharide (LPS). Acyltransferase activity is unchanged.,function:Possesses both acyltransferase and acetyltransferase activities. Activity is calcium-dependent. Involved in platelet-activating factor (PAF) biosynthesis by catalyzing the conversion of the PAF precursor, 1-O-alkyl-sn-glycero-3-phosphocholine (lyso-PAF) into 1-O-alkyl-2-acetyl-sn-glycero-3-phosphocholine (PAF). Also converts lyso-PAF to 1-alkyl-phosphatidylcholine (PC), a major component of cell membranes and a PAF precursor. Under resting conditions, acyltransferase activity is preferred. Upon acute inflammatory stimulus, acetyltransferase activity is enhanced and PAF synthesis increases.,pathway:Lipid metabolism; phospholipid metabolism.,similarity:Belongs to the 1-acyl-sn-glycerol-3-phosphate acyltransferase family.,similarity:Contains 2 EF-hand domains.,


Research Area

Image Data



Western blot analysis of lysates from K562 and HT-29 cells, using LPCAT2 Antibody. The lane on the right is blocked with the synthesized peptide.

Product Name: LPCAT2 Rabbit Polyclonal Antibody
Catalog #: APRab13389



Western blot analysis of the lysates from HT-29 cells using LPCAT2 antibody.

Note

For research use only.