

Summary

Production Name	MEK Kinase-3 Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human, Mouse

Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw
	cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	MAP3K3		
Alternative Names	MAP3K3; MAPKKK3; MEKK3; Mitogen-activated protein kinase kinase kinase 3;		
	MAPK/ERK kinase kinase 3; MEK kinase 3; MEKK 3		
Gene ID	4215.0		
SwissProt ID	Q99759. The antiserum was produced against synthesized peptide derived from human		
	MAP3K3. AA range:101-150		

Application

Dilution Ratio	WB 1:500 - 1:2000. ELISA: 1:20000
Molecular Weight	71kD



Background

This gene product is a 626-amino acid polypeptide that is 96.5% identical to mouse Mekk3. Its catalytic domain is closely related to those of several other kinases, including mouse Mekk2, tobacco NPK, and yeast Ste11. Northern blot analysis revealed a 4.6-kb transcript that appears to be ubiquitously expressed. This protein directly regulates the stress-activated protein kinase (SAPK) and extracellular signal-regulated protein kinase (ERK) pathways by activating SEK and MEK1/2 respectively; it does not regulate the p38 pathway. In cotransfection assays, it enhanced transcription from a nuclear factor kappa-B (NFKB)-dependent reporter gene, consistent with a role in the SAPK pathway. Alternatively spliced transcript variants encoding distinct isoforms have been observed. [provided by RefSeq, Jul 2008],catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,enzyme regulation:Activated by phosphorylation on Thr-530.,function:Component of a protein kinase signal transduction cascade. Mediates activation of the NF-kappa-B, AP1 and DDIT3 transcriptional regulators.,similarity:Belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family. MAP kinase kinase subfamily.,similarity:Contains 1 OPR domain.,similarity:Contains 1 protein kinase domain.,subunit:Binds both upstream activators and downstream substrates in multimolecular complexes. Part of a complex with MAP2K3, RAC1 and CCM2. Interacts with MAP2K5 and SPAG9.,

Research Area

Regulation of Actin Dynamics; SAPK_JNK; Cell Growth; Stem cell pathway; B Cell Receptor

Image Data



Western blot analysis of lysates from HepG2 cells, using MAP3K3 Antibody. The lane on the right is blocked with the synthesized peptide.





Western Blot analysis of various cells using MEK Kinase-3 Polyclonal Antibody diluted at 1: 2000

Note

For research use only.