
Summary

Production Name	MIG Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	IHC,ELISA
Reactivity	Human,Rat,Mouse

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	CXCL9 CMK MIG SCYB9
Alternative Names	C-X-C motif chemokine 9 (Gamma-interferon-induced monokine;Monokine induced by interferon-gamma;HuMIG;MIG;Small-inducible cytokine B9)
Gene ID	4283.0
SwissProt ID	Q07325.Synthetic peptide from human protein at AA range: 90-125

Application

Dilution Ratio	IHC-p 1:50-200, ELISA 1:10000-20000.
Molecular Weight	

Background

Product Name: MIG Rabbit Polyclonal Antibody
Catalog #: APRab13902




This antimicrobial gene encodes a protein thought to be involved in T cell trafficking. The encoded protein binds to C-X-C motif chemokine 3 and is a chemoattractant for lymphocytes but not for neutrophils. [provided by RefSeq, Sep 2014],function:Cytokine that affects the growth, movement, or activation state of cells that participate in immune and inflammatory response. Chemotactic for activated T-cells. Binds to CXCR3.,induction:By interferon gamma. The induction is enhanced by TNF-alpha in dermal fibroblasts and vein endothelial cells.,online information:CXCL9 entry,similarity:Belongs to the intercrine alpha (chemokine CxC) family.,


Research Area

Cytokine-cytokine receptor interaction;Chemokine;Toll_Like;

Image Data



Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:200

Note

For research use only.