

---

## Summary

<b>Production Name</b>	MVK Rabbit Polyclonal Antibody
<b>Description</b>	Rabbit Polyclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	IHC, WB, ELISA
<b>Reactivity</b>	Human, Monkey

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
<b>Purification</b>	Affinity purification

## Immunogen

<b>Gene Name</b>	MVK
<b>Alternative Names</b>	MVK; Mevalonate kinase; MK
<b>Gene ID</b>	4598.0
<b>SwissProt ID</b>	Q03426. The antiserum was produced against synthesized peptide derived from human Mevalonate Kinase. AA range: 151-200

## Application

<b>Dilution Ratio</b>	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:10000. Not yet tested in other applications.
<b>Molecular Weight</b>	42kD


## Background

This gene encodes the peroxisomal enzyme mevalonate kinase. Mevalonate is a key intermediate, and mevalonate kinase a key early enzyme, in isoprenoid and sterol synthesis. Mevalonate kinase deficiency caused by mutation of this gene results in mevalonic aciduria, a disease characterized psychomotor retardation, failure to thrive, hepatosplenomegaly, anemia and recurrent febrile crises. Defects in this gene also cause hyperimmunoglobulinaemia D and periodic fever syndrome, a disorder characterized by recurrent episodes of fever associated with lymphadenopathy, arthralgia, gastrointestinal distension and skin rash. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul 2014],catalytic activity:ATP + (R)-mevalonate = ADP + (R)-5-phosphomevalonate.,disease:Defects in MVK are the cause of hyperimmunoglobulinemia D and periodic fever syndrome (HIDS) [MIM:260920]. HIDS is an autosomal recessive disease characterized by recurrent episodes of unexplained high fever associated with skin rash, diarrhea, adenopathy (swollen, tender lymph nodes), arthralgias and/or arthritis. Concentration of IgD, and often IgA, are above normal.,disease:Defects in MVK are the cause of mevalonic aciduria [MIM:610377]. It is an accumulation of mevalonic acid which causes a variety of symptoms such as psychomotor retardation, dysmorphic features, cataracts, hepatosplenomegaly, lymphadenopathy, anemia, hypotonia, myopathy, and ataxia.,enzyme regulation:Farnesyl- and geranyl-pyrophosphates are competitive inhibitors.,function:May be a regulatory site in cholesterol biosynthetic pathway.,online information:Repertory of FMF and hereditary autoinflammatory disorders mutations,pathway:Isoprenoid biosynthesis; isopentenyl-PP biosynthesis via mevalonic acid pathway; isopentenyl-PP from (R)-mevalonic acid: step 1/3.,similarity:Belongs to the GHMP kinase family.,similarity:Belongs to the GHMP kinase family. Mevalonate kinase subfamily.,subunit:Homodimer.,

## Research Area


Terpenoid backbone biosynthesis;

## Image Data




Immunohistochemistry analysis of paraffin-embedded human brain tissue, using Mevalonate Kinase Antibody. The picture on the right is blocked with the synthesized peptide.

**Product Name: MVK Rabbit Polyclonal Antibody**  
**Catalog #: APRab14250**



Western blot analysis of lysates from COS7 and COLO205 cells, using Mevalonate Kinase Antibody. The lane on the right is blocked with the synthesized peptide.



Western Blot analysis of various cells using MVK Polyclonal Antibody

## Note

For research use only.