

## Summary

<b>Production Name</b>	NOXIN Rabbit Polyclonal Antibody
<b>Description</b>	Rabbit Polyclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB
<b>Reactivity</b>	Human,Rat,Mouse

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
<b>Purification</b>	Affinity purification

## Immunogen

<b>Gene Name</b>	NOXIN
<b>Alternative Names</b>	NOXIN; C11orf82; Nitric oxide-inducible gene protein
<b>Gene ID</b>	220042.0
<b>SwissProt ID</b>	Q8IXT1.The antiserum was produced against synthesized peptide derived from the Internal region of human NOXIN. AA range:111-160

## Application

<b>Dilution Ratio</b>	WB 1:500-1:2000. ELISA: 1:10000.
<b>Molecular Weight</b>	85kD

## Background

function:Acts as an anti-apoptotic factor and its absence increases cell death under normal and stress conditions. Can


**Product Name: NOXIN Rabbit Polyclonal Antibody**  
**Catalog #: APRab14820**



induce cell cycle arrest in the G1 or early S phase and this activity is independent of TP53/p53.,subcellular location:Accumulates in the nucleus in response to stress.,function:Acts as an anti-apoptotic factor and its absence increases cell death under normal and stress conditions. Can induce cell cycle arrest in the G1 or early S phase and this activity is independent of TP53/p53.,subcellular location:Accumulates in the nucleus in response to stress.,

## Research Area

## Image Data



Western Blot analysis of 823, 293T, AD293, Hela, HepG2 cells using NOXIN Polyclonal Antibody. Antibody was diluted at 1:2000. Secondary antibody was diluted at 1:20000

## Note

For research use only.