

# Summary

# Performance

Conjugation	Unconjugated	
Modification	Unmodified	
lsotype	IgG	
Clonality	Polyclonal	
Form	Liquid	
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw	
	cycles.	
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.	
Purification	Affinity purification	

# Immunogen

Gene Name	OR2H2
Alternative Names	
Gene ID	7932.0
SwissProt ID	O95918/Q2M1Y3/Q5SUK1.The antiserum was produced against synthesized peptide
	derived from human OR2H2. AA range:241-290

# Application

<b>Dilution Ratio</b>	WB 1:500 - 1:2000. ELISA: 1:10000
Molecular Weight	34kD

#### Background


olfactory receptor family 2 subfamily H member 2(OR2H2) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,

EnkiLife


### **Research Area**


Olfactory transduction;

# **Image Data**



Western blot analysis of lysates from Jurkat and MCF-7 cells, using OR2H2 Antibody. The lane on the right is blocked with







Note For research use only.