

Summary

tor 51D1 Rabbit Polyclonal Antibody
al Antibody
ise

Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw
	cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	OR51D1
Alternative Names	OR51D1; Olfactory receptor 51D1; Olfactory receptor OR11-14
Gene ID	390038.0
SwissProt ID	Q8NGF3.The antiserum was produced against synthesized peptide derived from
	human OR51D1. AA range:33-82

Application

Dilution Ratio	WB 1:500 - 1:2000. ELISA: 1:10000. Not yet tested in other applications.
Molecular Weight	34kD




Background

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,


Research Area

Olfactory transduction;

Image Data




Western blot analysis of lysates from Jurkat cells, using OR51D1 Antibody. The lane on the right is blocked with the



Western blot analysis of the lysates from COLO205 cells using OR51D1 antibody.





Western Blot analysis of various cells using Olfactory receptor 51D1 Polyclonal Antibody

Note

For research use only.