

Summary

Production Name	Palladin Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	IHC,ELISA
Reactivity	Human,Mouse

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	PALLD
Alternative Names	PALLD; KIAA0992; CGI-151; Palladin; SIH002; Sarcoma antigen NY-SAR-77
Gene ID	23022.0
SwissProt ID	Q8WX93.Synthesized peptide derived from Palladin . at AA range: 450-530

Application

Dilution Ratio	IHC 1:100-1:300 ELISA: 1:20000
Molecular Weight	150kD


Background

This gene encodes a cytoskeletal protein that is required for organizing the actin cytoskeleton. The protein is a component of actin-containing microfilaments, and it is involved in the control of cell shape, adhesion, and contraction. Polymorphisms

in this gene are associated with a susceptibility to pancreatic cancer type 1, and also with a risk for myocardial infarction. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Oct 2009],caution:Was wrongly assigned as myoneurin (Ref.2),disease:Genetic variations in PALLD are associated with susceptibility to pancreatic cancer type 1 (PNCA1) [MIM:606856]. Expression is increased early in the development of pancreatic cancer: in normal-appearing whole tissue immediately adjacent to cancer, in the pre-cancer, and in both the familial and sporadic forms of the cancer.,disease:Genetic variations in PALLD may be associated with myocardial infarction.,function:Cytoskeletal protein required for organization of normal actin cytoskeleton. Roles in establishing cell morphology, motility, cell adhesion and cell-extracellular matrix interactions in a variety of cell types. May function as a scaffolding molecule with the potential to influence both actin polymerization and the assembly of existing actin filaments into higher-order arrays. Binds to proteins that bind to either monomeric or filamentous actin. Localizes at sites where active actin remodeling takes place, such as lamellipodia and membrane ruffles. Different isoforms may have functional differences. Involved in the control of morphological and cytoskeletal changes associated with dendritic cell maturation. Involved in targeting ACTN to specific subcellular foci.,induction:Isoform 3 is expressed de novo and isoform 4 is up-regulated by TGF β 1 during myofibroblast differentiation.,miscellaneous:Protein is overexpressed in FA6, HPAF, IMIMPC2, SUIT2 and PATU2 sporadic pancreatic cancer cell lines.,PTM:Phosphorylated predominantly on serines and, to a lesser extent, on tyrosines (By similarity). Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the myotilin/palladin family.,similarity:Contains 5 Ig-like C2-type (immunoglobulin-like) domains.,subcellular location:Localizes to stress fibers and Z-disks. Cell junction, focal adhesion. Cell projection, ruffle. Cell projection, lamellipodium.,subunit:Interacts with EPS8, LASP1 and VASP (By similarity). Interacts with ACTN, ARGBP2, LPP, PFN1, SPIN90, SRC and EZR.,tissue specificity:Detected in both muscle and non-muscle tissues. High expression in prostate, ovary, colon, and kidney. Not detected in spleen, skeletal muscle, lung and peripheral blood lymphocytes (at protein level),.

Research Area

Image Data



Immunohistochemical analysis of paraffin-embedded human Colon cancer. 1, Antibody was diluted at 1:200 (4° overnight) .

Product Name: Palladin Rabbit Polyclonal Antibody
Catalog #: APRab15718



2, Tris-EDTA, pH 9.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200 (room temperature, 45min) .

Note

For research use only.