Catalog #: APRab15944



## **Summary**

| Production Name | PDZ-RhoGEF Rabbit Polyclonal Antibody |
|-----------------|---------------------------------------|
| Description     | Rabbit Polyclonal Antibody            |
| Host            | Rabbit                                |
| Application     | WB                                    |
| Reactivity      | Human, Mouse, Rat                     |

#### Performance

| Conjugation  | Unconjugated   |
|--------------|--|
| Modification | Unmodified   |
| lsotype      | lgG  |
| Clonality    | Polyclonal   |
| Form         | Liquid   |
| Storage      | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw   |
|              | cycles.  |
| Buffer       | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N. |
| Purification | Affinity purification  |

#### Immunogen

| Gene Name         | ARHGEF11   |
|-------------------|--|
| Alternative Names | ARHGEF11; KIAA0380; Rho guanine nucleotide exchange factor 11; PDZ-RhoGEF        |
| Gene ID           | 9826.0   |
| SwissProt ID      | O15085.The antiserum was produced against synthesized peptide derived from human |
|                   | ARHGEF11. AA range:198-247   |

# Application

| Dilution Ratio | WB 1:500-2000 |
|----------------|---------------|
|                |               |

**Molecular Weight** 


### Background

Rho GTPases play a fundamental role in numerous cellular processes that are initiated by extracellular stimuli that work through G protein coupled receptors. The encoded protein may form a complex with G proteins and stimulate Rhodependent signals. A similar protein in rat interacts with glutamate transporter EAAT4 and modulates its glutamate transport activity. Expression of the rat protein induces the reorganization of the actin cytoskeleton and its overexpression induces the formation of membrane ruffling and filopodia. Two alternative transcripts encoding different isoforms have been described. [provided by RefSeq, Jul 2008],domain:The poly-Pro region is essential for plasma membrane localization upon stimulation.,function:May play a role in the regulation of RhoA GTPase by guanine nucleotide-binding alpha-12 (GNA12) and alpha-13 (GNA13). Acts as guanine nucleotide exchange factor (GEF) for RhoA GTPase and may act as GTPase-activating protein (GAP) for GNA12 and GNA13.,similarity:Contains 1 DH (DBL-homology) domain.,similarity:Contains 1 PDZ (DHR) domain.,similarity:Contains 1 PH domain.,similarity:Contains 1 RGSL (RGS-like) domain.,subcellular location:Translocated to the membrane upon stimulation.,subunit:Interacts with GNA12 and GNA13 through the RGS domain. Interacts with RHOA, PLXNB1 and PLXNB2. Interacts with SLC1A6.,tissue specificity:Ubiquitously expressed.,


### **Research Area**

Regulation of Actin Dynamics; AMPK

### Image Data



Western blot analysis of ARHGEF11 Antibody. The lane on the right is blocked with the ARHGEF11 peptide.





Western Blot analysis of various cells using PDZ-RhoGEF Polyclonal Antibody

#### Note

For research use only.