

Product Name: PIASx Rabbit Polyclonal Antibody
Catalog #: APRab16121



Summary

Production Name	PIASx Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human,Mouse,Rat,Monkey

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	PIAS2
Alternative Names	PIAS2; PIASX; E3 SUMO-protein ligase PIAS2; Androgen receptor-interacting protein 3; ARIP3; DAB2-interacting protein; DIP; Msx-interacting zinc finger protein; Miz1; PIAS-NY protein; Protein inhibitor of activated STAT x; Protein inhibitor
Gene ID	9063.0
SwissProt ID	O75928.The antiserum was produced against synthesized peptide derived from human PIAS2. AA range:10-59

Application

Dilution Ratio	WB 1:500-2000; ELISA 2000-20000
Molecular Weight	68kD

Background


This gene encodes a member of the protein inhibitor of activated STAT (PIAS) family. PIAS proteins function as SUMO E3 ligases and play important roles in many cellular processes by mediating the sumoylation of target proteins. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. Isoforms of the encoded protein enhance the sumoylation of specific target proteins including the p53 tumor suppressor protein, c-Jun, and the androgen receptor. A pseudogene of this gene is located on the short arm of chromosome 4. The symbol MIZ1 has also been associated with ZBTB17 which is a different gene located on chromosome 1. [provided by RefSeq, Aug 2011],developmental stage:Expression of isoform 3 in adult testis is 14.2-fold stronger than in embryonic testis.,domain:The LXXLL motif is a transcriptional coregulator signature.,function:Functions as an E3-type small ubiquitin-like modifier (SUMO) ligase, stabilizing the interaction between UBE2I and the substrate, and as a SUMO-tethering factor. Plays a crucial role as a transcriptional coregulator in various cellular pathways, including the STAT pathway, the p53 pathway and the steroid hormone signaling pathway. The effects of this transcriptional coregulation, transactivation or silencing may vary depending upon the biological context and the PIAS2 isoform studied. However, it seems to be mostly involved in gene silencing. Binds to sumoylated ELK1 and enhances its transcriptional activity by preventing recruitment of HDAC2 by ELK1, thus reversing SUMO-mediated repression of ELK1 transactivation activity. Isoform PIAS2-beta, but not isoform PIAS2-alpha, promotes MDM2 sumoylation. Isoform PIAS2-alpha promotes PARK7 sumoylation. Isoform PIAS2-beta promotes NCOA2 sumoylation more efficiently than isoform PIAS2-alpha.,induction:Up-regulated transiently during myeloid differentiation in various cells lines, such as HL-60, U-937, K-562, induced by either phorbol ester (TPA) or retinoic acid.,pathway:Protein modification; protein sumoylation.,PTM:Sumoylated.,similarity:Belongs to the PIAS family.,similarity:Contains 1 SAP domain.,similarity:Contains 1 SP-RING-type zinc finger.,subcellular location:Colocalizes at least partially with promyelocytic leukemia nuclear bodies (PML NBs),subunit:Binds SUMO1 and UBE2I. Interacts with JUN, MDM2, PARK7, TP53 and TP73 isoform alpha, but not TP73 isoform beta. Interacts with STAT4 following IL12 and IFN-alpha stimulation of T-cells. Interacts also with GTF2I, GTF2IRD1, DAB2 and MSX2, as well as with several steroid receptors, including ESR1, ESR2, NR3C1, PGR, AR, and with NCOA2 (By similarity). Sumoylation of a target protein seems to enhance the interaction. Binds to sumoylated ELK1. Binds DNA, such as CDKN1A promoter, in a sequence-specific manner. Interacts with PLAG1. Interacts with KLF8; the interaction results in SUMO ligation and repression of KLF8 transcriptional activity and of its cell cycle progression into G(1) phase.,tissue specificity:Mainly testis. Isoform 3 is expressed predominantly in adult testis, weakly in pancreas, embryonic testis and sperm, and at very low levels in other organs.,

Research Area


Ubiquitin mediated proteolysis;Jak_STAT;Pathways in cancer;Small cell lung cancer;

Image Data


Product Name: PIASx Rabbit Polyclonal Antibody
Catalog #: APRab16121



Western blot analysis of lysates from COS7 cells, using PIAS2 Antibody. The lane on the right is blocked with the synthesized peptide.



Western Blot analysis of various cells using PIASx Polyclonal Antibody diluted at 1: 1000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Inventbiotech, MN, USA) .



Western Blot analysis of COS7 cells using PIASx Polyclonal Antibody diluted at 1: 1000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Inventbiotech, MN, USA) .

Note

For research use only.