# **Product Name: PIG3 Rabbit Polyclonal Antibody**

Catalog #: APRab16131



## **Summary**

**Production Name** PIG3 Rabbit Polyclonal Antibody

**Description** Rabbit Polyclonal Antibody

Host Rabbit
Application WB,ELISA

**Reactivity** Human, Rat, Mouse

#### **Performance**

ConjugationUnconjugatedModificationUnmodified

**Isotype** IgG

ClonalityPolyclonalFormLiquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw

cycles.

**Buffer** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.

**Purification** Affinity purification

#### **Immunogen**

Storage

Gene Name TP53I3

TP53I3; PIG3; Quinone oxidoreductase PIG3; Tumor protein p53-inducible protein 3; Alternative Names

p53-induced gene 3 protein

**Gene ID** 9540.0

Q53FA7.The antiserum was produced against synthesized peptide derived from human **SwissProt ID** 

QORX. AA range:241-290

## **Application**

**Dilution Ratio** WB 1:500 - 1:2000. ELISA: 1:20000

Molecular Weight 36kD

# **Product Name: PIG3 Rabbit Polyclonal Antibody**

Catalog #: APRab16131




## **Background**

The protein encoded by this gene is similar to oxidoreductases, which are enzymes involved in cellular responses to oxidative stresses and irradiation. This gene is induced by the tumor suppressor p53 and is thought to be involved in p53-mediated cell death. It contains a p53 consensus binding site in its promoter region and a downstream pentanucleotide microsatellite sequence. P53 has been shown to transcriptionally activate this gene by interacting with the downstream pentanucleotide microsatellite sequence. The microsatellite is polymorphic, with a varying number of pentanucleotide repeats directly correlated with the extent of transcriptional activation by p53. It has been suggested that the microsatellite polymorphism may be associated with differential susceptibility to cancer. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.alternative products:UV radiation favors the production of isoform 2,function:May be involved in the generation of reactive oxygen species (ROS),,induction:Isoforms 1 and 2 are both activated by TP53/p53, doxorubicin, etoposide and ionizing radiation. Isoform 2 is highly activated by UV radiation,,similarity:Belongs to the zinc-containing alcohol dehydrogenase family. Quinone oxidoreductase subfamily.


#### **Research Area**

p53;

## **Image Data**



Western blot analysis of lysates from 293 cells, using QORX Antibody. The lane on the right is blocked with the synthesized peptide.



# **Product Name: PIG3 Rabbit Polyclonal Antibody**

**C** EnkiLife

Catalog #: APRab16131

Western Blot analysis of various cells using PIG3 Polyclonal Antibody

#### Note

For research use only.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838