# **Product Name: PPP2R3C Rabbit Polyclonal Antibody**

Catalog #: APRab16429



### **Summary**

Production Name PPP2R3C Rabbit Polyclonal Antibody

**Description** Rabbit Polyclonal Antibody

Host Rabbit
Application WB,ELISA

**Reactivity** Human, Mouse, Rat

#### **Performance**

ConjugationUnconjugatedModificationUnmodified

**Isotype** IgG

ClonalityPolyclonalFormLiquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw

cycles.

**Buffer** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.

**Purification** Affinity purification

#### **Immunogen**

Storage

Gene Name PPP2R3C

PPP2R3C; C14orf10; G5PR; Serine/threonine-protein phosphatase 2A regulatory

Alternative Names subunit B" subunit gamma; Protein phosphatase subunit G5PR; Rhabdomyosarcoma

antigen MU-RMS-40.6A/6C

Gene ID 55012.0

Q969Q6.The antiserum was produced against synthesized peptide derived from human **SwissProt ID** 

PPP2R3C. AA range:165-214

## **Application**

**Dilution Ratio** WB 1:500-2000; ELISA 2000-20000

Molecular Weight 55kD

# **Product Name: PPP2R3C Rabbit Polyclonal Antibody**

Catalog #: APRab16429




### **Background**


This gene encodes a regulatory subunit of the serine/threonine phosphatase, protein phosphatase 2. This protein is localized to both nuclear and cytoplasmic regions depending on cell cycle phase. Homozygous conditional knockout mice for this gene exhibit reduced numbers and impaired proliferation of immune system B cells. This protein may regulate the expression of the P-glycoprotein ATP-binding cassette transporter through its phosphatase activity. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Feb 2015],developmental stage:Expressed in fetal brain.,function:May regulate MCM3AP phosphorylation through phosphatase recruitment. May play a role in the activation-induced cell death of B-cells,,similarity:Contains 2 EF-hand domains.,subcellular location:Excluded from the nucleoli. Localization is cell cycle-dependent. Localizes to the cytoplasm during cytokinesis,,subunit:Interacts with MCM3AP/GANP (By similarity). Interacts with PPP5C, and the phosphatase 2A core enzyme composed of the PPP2CA catalytic subunit and the constant regulatory subunit PPP2R1A,,tissue specificity:Ubiquitously expressed in brain and other tissues.,

#### Research Area

#### **Image Data**




Western blot analysis of PPP2R3C Antibody. The lane on the right is blocked with the PPP2R3C peptide.



Western blot analysis of the lysates from HepG2 cells using PPP2R3C antibody.

# Product Name: PPP2R3C Rabbit Polyclonal Antibody Catalog #: APRab16429





Western Blot analysis of various cells using PPP2R3C Polyclonal Antibody

#### Note

For research use only.