

Summary

Production Name	Septin 3 Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human, Mouse, Rat

Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	lgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw
	cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	SEPT3
Alternative Names	SEPT3; SEP3; Neuronal-specific septin-3
Gene ID	55964.0
SwissProt ID	Q9UH03.The antiserum was produced against synthesized peptide derived from
	human SEPT3. AA range:136-185

Application

Dilution Ratio	WB 1:500-2000; ELISA 2000-20000
Molecular Weight	41kD

Background


Product Name: Septin 3 Rabbit Polyclonal Antibody Catalog #: APRab17741




septin 3(SEPT3) Homo sapiens This gene belongs to the septin family of GTPases. Members of this family are required for cytokinesis. Expression is upregulated by retinoic acid in a human teratocarcinoma cell line. The specific function of this gene has not been determined. Alternative splicing of this gene results in two transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008],function:Involved in cytokinesis .,induction:Up-regulated during neuronal differentiation.,PTM:Phosphorylated by PKG on serine residues. Phosphorylated by PKG on Ser-91.,similarity:Belongs to the septin family.,subcellular location:Localizes in synaptosomes at dendritic synapses of neurons. Localizes within presynaptic terminals of neurons.,subunit:May assemble into a multicomponent structure.,tissue specificity:Brain specific.,

Research Area

Image Data




Western blot analysis of lysates from 293 cells, using SEPT3 Antibody. The lane on the right is blocked with the synthesized



Western Blot analysis of various cells using Septin 3 Polyclonal Antibody diluted at 1: 1000





Western blot analysis of various lysis using Septin 3 Polyclonal Antibody diluted at 1: 1000. Secondary antibody was diluted at 1:20000

Note

For research use only.