

# Summary

Production Name	SGLT-2 Rabbit Polyclonal Antibody	
Description	Rabbit Polyclonal Antibody	
Host	Rabbit	
Application	WB	
Reactivity	Human, Mouse, Rat	

#### Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw
	cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

## Immunogen

Gene Name	SLC5A2
Alternative Names	SLC5A2; SGLT2; Sodium/glucose cotransporter 2; Na(+)/glucose cotransporter 2; Low
	affinity sodium-glucose cotransporter; Solute carrier family 5 member 2
Gene ID	6524.0
SwissProt ID	P31639.The antiserum was produced against synthesized peptide derived from human
	SLC5A2. AA range:101-150

# Application

Dilution Ratio	WB 1:500-2000
Molecular Weight	73kD



### Background

This gene encodes a member of the sodium glucose cotransporter family which are sodium-dependent glucose transport proteins. The encoded protein is the major cotransporter involved in glucose reabsorption in the kidney. Mutations in this gene are associated with renal glucosuria. Two transcript variants, one protein-coding and one not, have been found for this gene. [provided by RefSeq, Feb 2015],disease:Defects in SLC5A2 are the cause of renal glucosuria (GLYS1) [MIM:233100]. GLYS1 is an autosomal recessive disorder characterized by a normal fasting serum glucose concentration and persistent isolated glucosuria, with a normal glucose tolerance test.,function:Efficient substrate transport in mammalian kidney is provided by the concerted action of a low affinity high capacity and a high affinity low capacity Na(+)/glucose cotransporter arranged in series along kidney proximal tubules.,function:Sodium-dependent glucose transporter. Has a Na(+) to glucose coupling ratio of 1:1.,similarity:Belongs to the sodium:solute symporter (SSF) (TC 2.A.21) family.,

### **Research Area**

Image Data

### 

Western blot analysis of lysates from 293 cells, using SLC5A2 Antibody. The lane on the right is blocked with the synthesized







Note

For research use only.