

**Product Name: SP-100 Rabbit Polyclonal Antibody**  
**Catalog #: APRab18147**



## Summary

<b>Production Name</b>	SP-100 Rabbit Polyclonal Antibody
<b>Description</b>	Rabbit Polyclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	IHC,ELISA
<b>Reactivity</b>	Human,Rat,Mouse

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
<b>Purification</b>	Affinity purification

## Immunogen

<b>Gene Name</b>	SP100
<b>Alternative Names</b>	SP100; Nuclear autoantigen Sp-100; Lysp100b; Nuclear dot-associated Sp100 protein; Speckled 100 kDa
<b>Gene ID</b>	6672.0
<b>SwissProt ID</b>	P23497.Synthesized peptide derived from SP-100 . at AA range: 250-330

## Application

<b>Dilution Ratio</b>	IHC 1:100-1:300 ELISA: 1:40000
<b>Molecular Weight</b>	100kD


## Background

This gene encodes a subnuclear organelle and major component of the PML (promyelocytic leukemia)-SP100 nuclear

bodies. PML and SP100 are covalently modified by the SUMO-1 modifier, which is considered crucial to nuclear body interactions. The encoded protein binds heterochromatin proteins and is thought to play a role in tumorigenesis, immunity, and gene regulation. Alternatively spliced variants have been identified for this gene; one of which encodes a high-mobility group protein. [provided by RefSeq, Aug 2011],disease:This antigen is recognized by autoantibodies from patients with primary biliary cirrhosis (PBC),.domain:Contains one Pro-Xaa-Val-Xaa-Leu (PxVxL) motif, which is required for interaction with chromoshadow domains. This motif requires additional residues -7, -6, +4 and +5 of the central Val which contact the chromoshadow domain,.domain:The HSR domain is important for the nuclear body targeting as well as for the dimerization,.function:May play a role in the control of gene expression,.induction:By interferon,.miscellaneous:The major isoform Sp100-A, has a calculated MW of 54 kDa, but exhibits aberrant electrophoretic mobilities, with an apparent MW OF 100 kDa,.PTM:Phosphorylated,.PTM:Sumoylated. Sumoylation depends on a functional nuclear localization signal but is not necessary for nuclear import or nuclear body targeting,.similarity:Contains 1 HSR domain,.similarity:Contains 1 SAND domain,.similarity:Contains 2 HMG box DNA-binding domains,.subcellular location:Found in the nuclear body, also known as nuclear domain 10 (ND10), PML oncogenic domain (POD), nuclear dots (ND) and KR body. The nuclear body is a nucleoplasmic structure of punctate shape, which varies in size and number. Induction by interferon and may be cell cycle stages modulate the subnuclear localization of the isoforms,.subunit:Homodimer. Splice variants heterodimerize. Interacts with members of the HP1 family of nonhistone chromosomal protein, such as CBX5 and CBX3 via the PxVxL motif. Interacts with Epstein-Barr virus EBNA-LP,.tissue specificity:Widely expressed. Sp100-B is expressed only in spleen, tonsil, thymus, mature B-cell line and some T-cell line, but not in brain, liver, muscle or non-lymphoid cell lines,.

## Research Area

## Image Data



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200 (4° overnight) . 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 45min) .

**Product Name: SP-100 Rabbit Polyclonal Antibody**  
**Catalog #: APRab18147**



---

**Note**

For research use only.