

---

## Summary

<b>Production Name</b>	SPOP Rabbit Polyclonal Antibody
<b>Description</b>	Rabbit Polyclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB,IHC,
<b>Reactivity</b>	Human,Mouse,Rat

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
<b>Purification</b>	Affinity purification

## Immunogen

<b>Gene Name</b>	SPOP
<b>Alternative Names</b>	Speckle-type POZ protein (HIB homolog 1) (Roadkill homolog 1)
<b>Gene ID</b>	8405.0
<b>SwissProt ID</b>	O43791.The antiserum was produced against synthesized peptide derived from the Internal region of human SPOP. AA range:41-90

## Application

<b>Dilution Ratio</b>	WB 1:500-2000;IHC 1:50-300
<b>Molecular Weight</b>	42kD

## Background

---


**Product Name: SPOP Rabbit Polyclonal Antibody**  
**Catalog #: APRab18207**



This gene encodes a protein that may modulate the transcriptional repression activities of death-associated protein 6 (DAXX), which interacts with histone deacetylase, core histones, and other histone-associated proteins. In mouse, the encoded protein binds to the putative leucine zipper domain of macroH2A1.2, a variant H2A histone that is enriched on inactivated X chromosomes. The BTB/POZ domain of this protein has been shown in other proteins to mediate transcriptional repression and to interact with components of histone deacetylase co-repressor complexes. Alternative splicing of this gene results in multiple transcript variants encoding the same protein. [provided by RefSeq, Jul 2008],domain:The MATH domain interacts with H2AFY and BMI1.,function:Inhibits IPF1/PDX1 transactivation of established target promoters, such as insulin, may be by recruiting a repressor complex (By similarity). In complex with CUL3, involved in ubiquitination of BMI1, H2AFY and DAXX, and probably also in ubiquitination and proteasomal degradation of Gli2 or Gli3.,miscellaneous:Antigen recognized by serum from scleroderma patient.,pathway:Protein modification; protein ubiquitination.,similarity:Belongs to the Tdpoz family.,similarity:Contains 1 BTB (POZ) domain.,similarity:Contains 1 MATH domain.,subunit:Homodimer. Part of a complex consisting of BMI1, CUL3 and SPOP. Part of a complex consisting of H2AFY, CUL3 and SPOP. Part of a complex consisting of DAXX, CUL3 and SPOP. Interacts with H2AFY, IPF1/PDX1, BMI1 and DAXX. Interacts with CUL3.,tissue specificity:Widely expressed.,


## Research Area

## Image Data




Western Blot analysis of 3T3 HEPG2 HeLa cells using SPOP Polyclonal Antibody diluted at 1:1500. Secondary antibody was diluted at 1:20000


**Product Name: SPOP Rabbit Polyclonal Antibody**  
**Catalog #: APRab18207**




Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:200



**Product Name: SPOP Rabbit Polyclonal Antibody**  
**Catalog #: APRab18207**



---

Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:200

**Note**

For research use only.