

Summary

| Production Name | T2R49 Rabbit Polyclonal Antibody |
|-----------------|----------------------------------|
| Description | Rabbit Polyclonal Antibody |
| Host | Rabbit |
| Application | WB,ELISA |
| Reactivity | Human,Rat,Mouse |

Performance

| Conjugation | Unconjugated |
|--------------|--|
| Modification | Unmodified |
| lsotype | IgG |
| Clonality | Polyclonal |
| Form | Liquid |
| Storage | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw |
| | cycles. |
| Buffer | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N. |
| Purification | Affinity purification |

Immunogen

| Gene Name | TAS2R20 |
|-------------------|--|
| Alternative Names | TAS2R20; TAS2R49; Taste receptor type 2 member 20; Taste receptor type 2 member |
| | 49; T2R49; Taste receptor type 2 member 56; T2R56 |
| Gene ID | 259295.0 |
| SwissProt ID | P59543.The antiserum was produced against synthesized peptide derived from human |
| | TAS2R49. AA range:94-143 |

Application

| Dilution Ratio | WB 1:500 - 1:2000. ELISA: 1:20000 |
|------------------|-----------------------------------|
| Molecular Weight | 38kD |



Background

This gene encodes a member of the taste receptor two family of class C G-protein coupled receptors. Receptors of this family have a short extracellular N-terminus, seven transmembrane helices, three extracellular loops and three intracellular loops, and an intracellular C-terminus. Members of this family are expressed in a subset of taste receptor cells, where they function in bitter taste reception, as well as in non-gustatory cells including those of the brain, reproductive organs, respiratory system, and gastrointestinal system. This gene maps to the taste receptor gene cluster on chromosome 12p13. [provided by RefSeq, Jul 2016],function:Receptor that may play a role in the perception of bitterness and is gustducin-linked. May play a role in sensing the chemical composition of the gastrointestinal content. The activity of this receptor may stimulate alpha gustducin, mediate PLC-beta-2 activation and lead to the gating of TRPM5.,miscellaneous:Most taste cells may be activated by a limited number of bitter compounds; individual taste cells can discriminate among bitter stimuli.,similarity:Belongs to the G-protein coupled receptor T2R family.,tissue specificity:Expressed in subsets of taste receptor cells of the tongue and exclusively in gustducin-positive cells.,

Research Area

Taste transduction;

Image Data



Western blot analysis of lysates from COLO and HepG2 cells, using TAS2R49 Antibody. The lane on the right is blocked with

the synthesized peptide. (kD) 117-85-48-34-26-19-

Product Name: T2R49 Rabbit Polyclonal Antibody Catalog #: APRab18574



Western blot analysis of the lysates from COLO205 cells using TAS2R49 antibody.

Note

For research use only.