

Summary

Production Name	TAF II p250 Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	IHC,ELISA
Reactivity	Human,Mouse

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	TAF1
Alternative Names	TAF1; BA2R; CCG1; CCGS; TAF2A; Transcription initiation factor TFIID subunit 1; Cell cycle gene 1 protein; TBP-associated factor 250 kDa; p250; Transcription initiation factor TFIID 250 kDa subunit; TAF(II)250; TAFII-250; TAFII250
Gene ID	6872.0
SwissProt ID	P21675.The antiserum was produced against synthesized peptide derived from human TAF1. AA range:1131-1180

Application

Dilution Ratio	IHC 1:100-1:300 ELISA: 1:10000
Molecular Weight	

Background


Initiation of transcription by RNA polymerase II requires the activities of more than 70 polypeptides. The protein that coordinates these activities is the basal transcription factor TFIID, which binds to the core promoter to position the polymerase properly, serves as the scaffold for assembly of the remainder of the transcription complex, and acts as a channel for regulatory signals. TFIID is composed of the TATA-binding protein (TBP) and a group of evolutionarily conserved proteins known as TBP-associated factors or TAFs. TAFs may participate in basal transcription, serve as coactivators, function in promoter recognition or modify general transcription factors (GTFs) to facilitate complex assembly and transcription initiation. This gene encodes the largest subunit of TFIID. This subunit binds to core promoter sequences encompassing the transcription start site. It also bincatalytic activity:ATP + a phosphoprotein.,cofactor:Magnesium.,disease:Defects in TAF1 are the cause of dystonia type 3 (DYT3) [MIM:314250]; also called X-linked dystonia-parkinsonism (XDP). DYT3 is a X-linked dystonia-parkinsonism disorder. Dystonia is defined by the presence of sustained involuntary muscle contractions, often leading to abnormal postures. DYT3 is characterized by severe progressive torsion dystonia followed by parkinsonism. Its prevalence is high in the Philippines. DYT3 has a well-defined pathology of extensive neuronal loss and mosaic gliosis in the striatum (caudate nucleus and putamen) which appears to resemble that in Huntington disease.,enzyme regulation:Autophosphorylates on Ser residues. Inhibited by retinoblastoma tumor suppressor protein, RB1.,function:Largest component and core scaffold of the TFIID basal transcription factor complex. Contains novel N- and C-terminal Ser/Thr kinase domains which can autophosphorylate or transphosphorylate other transcription factors. Phosphorylates TP53 on 'Thr-55' which leads to MDM2-mediated degradation of TP53. Phosphorylates GTF2A1 and GTF2F1 on Ser residues. Possesses DNA-binding activity. Essential for progression of the G1 phase of the cell cycle.,PTM:Phosphorylated by casein kinase II in vitro.,similarity:Belongs to the TAF1 family.,similarity:Contains 1 HMG box DNA-binding domain.,similarity:Contains 2 bromo domains.,similarity:Contains 2 protein kinase domains.,subunit:TAF1 is the largest component of transcription factor TFIID that is composed of TBP and a variety of TBP-associated factors. TAF1, when part of the TFIID complex, interacts with C-terminus of TP53. RB1 interacts with the N-terminal domain of TAF1. Interacts with ASF1A and ASF1B. Interacts with SV40 Large T antigen.,

Research Area

Protein_Acetylation

Image Data

Product Name: TAF II p250 Rabbit Polyclonal Antibody
Catalog #: APRab18611



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using TAF1 Antibody. The picture on the right is blocked with the synthesized peptide.

Note

For research use only.