

---

## Summary

<b>Production Name</b>	Thrombin R Rabbit Polyclonal Antibody
<b>Description</b>	Rabbit Polyclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB,ELISA
<b>Reactivity</b>	Human,Rat,Mouse

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
<b>Purification</b>	Affinity purification

## Immunogen

<b>Gene Name</b>	F2R
<b>Alternative Names</b>	F2R; CF2R; PAR1; TR; Proteinase-activated receptor 1; PAR-1; Coagulation factor II receptor; Thrombin receptor
<b>Gene ID</b>	2149.0
<b>SwissProt ID</b>	P25116.The antiserum was produced against synthesized peptide derived from human Thrombin Receptor. AA range:10-59

## Application

<b>Dilution Ratio</b>	WB 1:500 - 1:2000. ELISA: 1:5000.
<b>Molecular Weight</b>	48kD


## Background

Coagulation factor II receptor is a 7-transmembrane receptor involved in the regulation of thrombotic response. Proteolytic cleavage leads to the activation of the receptor. F2R is a G-protein coupled receptor family member. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Aug 2015],function:High affinity receptor for activated thrombin coupled to G proteins that stimulate phosphoinositide hydrolysis. May play a role in platelets activation and in vascular development.,PTM:A proteolytic cleavage generates a new N-terminus that functions as a tethered ligand.,PTM:Phosphorylated; probably mediating desensitization prior to the uncoupling and internalization of the receptor.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Platelets and vascular endothelial cells.,

## Research Area


Akt\_PKB;Calcium;Neuroactive ligand-receptor interaction;Endocytosis;Complement and coagulation cascades;Regulates Actin and Cytoskeleton;

## Image Data



Western blot analysis of lysates from HeLa cells, treated with Nocodazole 1ug/ml 18h, using Thrombin Receptor Antibody.

The lane on the right is blocked with the synthesized peptide.



Western Blot analysis of various cells using Thrombin R Polyclonal Antibody

**Product Name: Thrombin R Rabbit Polyclonal Antibody**  
**Catalog #: APRab18889**



---

**Note**

For research use only.