
Summary

Production Name	TNF-IP 1 Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	TNFAIP1 TNFAIP1; BACURD2; EDP1; BTB/POZ domain-containing adapter for CUL3-mediated
Alternative Names	RhoA degradation protein 2; hBACURD2; BTB/POZ domain-containing protein TNFAIP1; Protein B12; Tumor necrosis factor; alpha-induced protein 1, endothelial
Gene ID	7126.0
SwissProt ID	Q13829.The antiserum was produced against synthesized peptide derived from human TNAP1. AA range:71-120

Application


Dilution Ratio	WB 1:500 - 1:2000. ELISA: 1:5000.
Molecular Weight	34kD

Background


This gene was identified as a gene whose expression can be induced by the tumor necrosis factor alpha (TNF) in umbilical vein endothelial cells. Studies of a similar gene in mouse suggest that the expression of this gene is developmentally regulated in a tissue-specific manner. [provided by RefSeq, Jul 2008],function:Enhances the PCNA-dependent DNA polymerase delta activity.,induction:By TNF-alpha, interleukin-1 beta and lipopolysaccharide (LPS).,similarity:Belongs to the KCTD10/KCTD13/TNFAIP1 family.,similarity:Contains 1 BTB (POZ) domain.,subunit:Interacts with PCNA.,

Research Area

Image Data




Western blot analysis of lysates from HUVEC cells, treated with PMA 125ng/ml 30', using TNAP1 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from 293 cells using TNAP1 antibody.

Product Name: TNF-IP 1 Rabbit Polyclonal Antibody
Catalog #: APRab19089



Western Blot analysis of various cells using TNF-IP 1 Polyclonal Antibody diluted at 1 : 500. Secondary antibody was diluted at 1:20000

Note

For research use only.