

**Product Name: USP48 Rabbit Polyclonal Antibody**  
**Catalog #: APRab19684**



## Summary

<b>Production Name</b>	USP48 Rabbit Polyclonal Antibody
<b>Description</b>	Rabbit Polyclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	IHC,ELISA
<b>Reactivity</b>	Human,Mouse,Rat

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
<b>Purification</b>	Affinity purification

## Immunogen

<b>Gene Name</b>	USP48
<b>Alternative Names</b>	USP48; USP31; Ubiquitin carboxyl-terminal hydrolase 48; Deubiquitinating enzyme 48; Ubiquitin thioesterase 48; Ubiquitin-specific-processing protease 48
<b>Gene ID</b>	84196.0
<b>SwissProt ID</b>	Q86UV5.The antiserum was produced against synthesized peptide derived from human USP48. AA range:631-680

## Application

<b>Dilution Ratio</b>	IHC 1:100-1:300 ELISA: 1:5000
<b>Molecular Weight</b>	120kD

## Background


**Product Name: USP48 Rabbit Polyclonal Antibody**  
**Catalog #: APRab19684**



This gene encodes a protein containing domains that associate it with the peptidase family C19, also known as family 2 of ubiquitin carboxyl-terminal hydrolases. Family members function as deubiquitinating enzymes, recognizing and hydrolyzing the peptide bond at the C-terminal glycine of ubiquitin. Enzymes in peptidase family C19 are involved in the processing of poly-ubiquitin precursors as well as that of ubiquitinated proteins. Alternate transcriptional splice variants, encoding different isoforms, have been characterized. [provided by RefSeq, Jul 2008],catalytic activity:Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol.,caution:Was named USP31 by some authors (PubMed:15354349 and PubMed:17081983),.function:Recognizes and hydrolyzes the peptide bond at the C-terminal Gly of ubiquitin. Involved in the processing of poly-ubiquitin precursors as well as that of ubiquitinated proteins. May be involved in the regulation of NF-kappa-B activation by TNF receptor superfamily via its interactions with RELA and TRAF2. May also play a regulatory role at post-synaptic sites.,sequence caution:Contaminating sequence. Potential poly-A sequence.,similarity:Belongs to the peptidase C19 family.,similarity:Contains 1 ubiquitin-like domain.,similarity:Contains 3 DUSP domains.,subunit:Interacts with TRAF2 and RELA.,tissue specificity:Widely expressed.,

## Research Area

## Image Data



Immunohistochemistry analysis of paraffin-embedded human colon carcinoma tissue, using USP48 Antibody. The picture on the right is blocked with the synthesized peptide.

## Note

For research use only.