

Product Name: β -1,4-GalNAc-T Rabbit Polyclonal Antibody
Catalog #: APRab20342

Summary

Production Name	β -1,4-GalNAc-T Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	IHC,ELISA
Reactivity	Human,Mouse,Rat

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	CSGALNACT1
Alternative Names	CSGALNACT1; CHGN; GALNACT1; Chondroitin sulfate N-acetylgalactosaminyltransferase 1; CsGalNAcT-1; Chondroitin beta-1; 4-N-acetylgalactosaminyltransferase 1; Beta4GalNAcT-1
Gene ID	55790.0
SwissProt ID	Q8TDX6.The antiserum was produced against synthesized peptide derived from human CSGALNACT1. AA range:201-250

Application

Dilution Ratio	IHC 1:100-1:300 ELISA: 1:40000
Molecular Weight	60kD

Product Name: β -1,4-GalNAc-T Rabbit Polyclonal Antibody
Catalog #: APRab20342




Background

catalytic activity:UDP-N-acetyl-D-galactosamine + beta-D-glucuronyl-(1->3)-D-galactosyl-proteoglycan = UDP + N-acetyl-D-galactosaminy-(1->4)-beta-D-glucuronyl-(1->3)-beta-D-galactosylproteoglycan.,function:Transfers 1,4-N-acetylgalactosamine (GalNAc) from UDP-GalNAc to the non-reducing end of glucuronic acid (GlcUA). Required for addition of the first GalNAc to the core tetrasaccharide linker and for elongation of chondroitin chains. Important role in chondroitin chain biosynthesis in cartilage.,online information:Chondroitin beta-1,4-N-acetylgalactosaminyltransferase 1,online information:GlycoGene database,PTM:N-glycosylated.,similarity:Belongs to the chondroitin N-acetylgalactosaminyltransferase family.,tissue specificity:Ubiquitous, with the highest levels in placenta, thyroid, bladder, prostate and adrenal gland. Detected at low levels in the other tissues examined.,catalytic activity:UDP-N-acetyl-D-galactosamine + beta-D-glucuronyl-(1->3)-D-galactosyl-proteoglycan = UDP + N-acetyl-D-galactosaminy-(1->4)-beta-D-glucuronyl-(1->3)-beta-D-galactosylproteoglycan.,function:Transfers 1,4-N-acetylgalactosamine (GalNAc) from UDP-GalNAc to the non-reducing end of glucuronic acid (GlcUA). Required for addition of the first GalNAc to the core tetrasaccharide linker and for elongation of chondroitin chains. Important role in chondroitin chain biosynthesis in cartilage.,online information:Chondroitin beta-1,4-N-acetylgalactosaminyltransferase 1,online information:GlycoGene database,PTM:N-glycosylated.,similarity:Belongs to the chondroitin N-acetylgalactosaminyltransferase family.,tissue specificity:Ubiquitous, with the highest levels in placenta, thyroid, bladder, prostate and adrenal gland. Detected at low levels in the other tissues examined.,

Research Area

Chondroitin sulfate biosynthesis;

Image Data



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using CSGALNACT1 Antibody. The picture on the right is blocked with the synthesized peptide.

**Product Name: β -1,4-GalNAc-T Rabbit Polyclonal
Antibody**
Catalog #: APRab20342



Note

For research use only.