


Summary

Name	R-Spondin1
Purity	Greater than 98% as determined by reducing SDS-PAGE
Endotoxin level	≤10 EU/mg
Construction	Recombinant Human R-Spondin1 is produced by our Mammalian cell expression system and the target gene encoding Ser21-Ala263 is expressed.
Accession #	Q2MKA7
Host	Human Cells
Species	Human
Predicted Molecular Mass	26.7 kDa
- 1.4	
Formulation	Lyophilized From PBS,5% mannitol and 0.01% Tween 80, pH7.4
Formulation	Lyophilized From PBS,5% mannitol and 0.01% Tween 80, pH7.4 The product is shipped on dry ice/polar packs.Upon receipt, store it immediately at the temperature listed below.
	The product is shipped on dry ice/polar packs.Upon receipt, store it immediately

SDS-PAGE image



Background



Alternative NamesRSPO1; R-spondin1; RP11-566C13.1; CRISTIN3; FLJ40906; RSPO Rspo1; R-spondin;
Rspondin; RP23-325M14.2; Roof plate-specific spondin-1BackgroundRSPO1 is a secreted protein, containing 2 FU(furin-like) repeats and 1 TSP type-1
domain and belonging to the R-spondin family. RSPO1 is required for the early
development of gonads, regardless of sex. It has been found in mice only eleven
days after fertilization. To induce cell proliferation, it acts synergistically with
WNT4. They help stabilize β catenin, which activates downstream targets. RSPO1 is
necessary in female sex development. It augments the WNT/β catenin pathway to
oppose male sex development. In critical gonadal stages, between six and nine
weeks after fertilization, the ovaries upregulate it while the testes downregulate it.
RSPO1 can potentially aid in the treatment of mucositis, which is characterized by
inflammation of the oral cavity. This unfortunate condition often accompanies
chemotherapy and radiation in cancer patients with head and neck tumors.

Note

For Research Use Only, Not for Diagnostic Use.