


Summary

Name	SDGF/Amphiregulin/AREG/AREGB/CRDGF/AR
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/ μ g as determined by LAL test.
Construction	Recombinant Human Amphiregulin is produced by our E.coli expression system and the target gene encoding Ser101-Lys198 is expressed.
Accession #	P15514
Host	E.coli
Species	Human
Predicted Molecular Mass	11.4 KDa
Formulation	Lyophilized from a 0.2 μ m filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at $\leq -20^{\circ}\text{C}$, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8 $^{\circ}\text{C}$ for 2-7 days. Aliquots of reconstituted samples are stable at $\leq -20^{\circ}\text{C}$ for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human AREG
Catalog #: PEH0058

Alternative Names

Amphiregulin; AR; Colorectum Cell-Derived Growth Factor; CRDGF; AREG; SDGF; AREGB

Background

Amphiregulin (AREG) is a single-pass membrane protein with 252 amino acids. AREG belongs to the amphiregulin family, which contains 1 EGF-like domain. AREG is expressed in a variety of tissues including ovary, placenta, lung, kidney, stomach, colon, and breast. It is related to Epidermal Growth Factor (EGF) and Transforming Growth Factor Alpha (TGF-alpha). As an EGF-related growth factor, AREG interacts with the EGF/TGF-alpha receptor to promote the growth of normal epithelial cells and inhibits the growth of certain aggressive carcinoma cell lines. AREG may also play a protective role in Bleomycin-Induced Pneumopathy.

Note

For Research Use Only , Not for Diagnostic Use.