


Summary

Name	ARC/NOL3
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Nucleolar Protein 3 is produced by our E.coli expression system and the target gene encoding Met1-Ser208 is expressed with a GST tag at the N-terminus.
Accession #	O60936
Host	E.coli
Species	Human
Predicted Molecular Mass	48.92 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM Tris-HCl, 350mM NaCl, 1mM DTT, pH 8.0.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human NOL3 (N-GST)
Catalog #: PEH0095



Alternative Names

Nucleolar Protein 3; Apoptosis Repressor With CARD; Muscle-Enriched Cytoplasmic Protein; Myp; Nucleolar Protein of 30 kDa; Nop30; NOL3; ARC; NOP

Background

Nucleolar Protein 3 is encoded by NOL3 gene; multiple transcript variants encoding different isoforms have been found for this gene. So far, Nucleolar protein 3 has show to have two Isoforms. Isoform 1 may be involved in RNA splicing.Isoform 2 may inhibit apoptosis.It has been shown to down-regulate the enzyme activities of caspase 2, caspase 8 and tumor protein p53.

Note

For Research Use Only , Not for Diagnostic Use.