


## Summary

<b>Name</b>	CD40 Ligand/CD40L/TNFSF5/CD154/T-cell antigen Gp39/gp39/TNF-related activation protein/TRAP/Tumor Necrosis Factor Ligand Superfamily Member 5/CD40LG
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE
<b>Endotoxin level</b>	<1 EU/ $\mu$ g as determined by LAL test.
<b>Construction</b>	Recombinant Human CD40L is produced by our E.coli expression system and the target gene encoding Met113-Leu261 is expressed.
<b>Accession #</b>	P29965
<b>Host</b>	E.coli
<b>Species</b>	Human
<b>Predicted Molecular Mass</b>	16.2 KDa
<b>Formulation</b>	Lyophilized from a 0.2 $\mu$ m filtered solution of 20mM PB, 200mM NaCl, 0.1mM EDTA, pH 7.0.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
<b>Stability&amp;Storage</b>	Store at $\leq$ -70°C, stable for 6 months after receipt. Store at $\leq$ -70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
<b>Reconstitution</b>	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 $\mu$ g/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## SDS-PAGE image



## Background

**Product Name: Recombinant Human CD40L**  
**Catalog #: PEH0347**

---

**Alternative Names**

CD40 Ligand; CD40-L; T-Cell Antigen Gp39; TNF-Related Activation Protein; TRAP; Tumor Necrosis Factor Ligand Superfamily Member 5; CD154; CD40LG; CD40L; TNFSF5; TRAP

**Background**

CD40 Ligand (CD40LG) is a type II transmembrane glycoprotein that belongs to the TNF superfamily. Like other TNF superfamily members, CD40LG exists as a trimer in membrane bound and soluble form, both of which are bioactive. CD40LG is a ligand for CD40; its ligation also initiates signal transduction in CD40LG expressing cells. CD40LG is a differentiation antigen that is expressed on the surface of T-cells. It stimulates B-cell proliferation and secretion of all immunoglobulin isotypes in the presence of cytokines. CD40LG has been shown to induce cytokine production and tumoricidal activity in peripheral blood monocytes. It also co-stimulates proliferation of activated T-cells and this is accompanied by the production of IFN-gamma, TNF-alpha, and IL2.

**Note**

For Research Use Only , Not for Diagnostic Use.