


## Summary

| Name                     | Cystathionine gamma-lyase/CTH  |
|--------------------------|--|
| Purity                   | Greater than 95% as determined by reducing SDS-PAGE  |
| Endotoxin level          | <1 EU/µg as determined by LAL test.  |
| Construction Accession # | Recombinant Human Cystathionine Gamma-Lyase is produced by our E.coli expression system and the target gene encoding Met1-Ser405 is expressed.<br>P32929                         |
|                          |  |
| Host                     | E.coli   |
| Species                  | Human  |
| Predicted Molecular Mass | 44.7 KDa   |
| Formulation              | Supplied as a 0.2 $\mu m$ filtered solution of 20mM Tris-HCl, 8% Sucrose, 0.05% Tween 80, pH8.0.   |
| Shipping                 | The product is shipped on dry ice/polar packs. Upon receipt, store it immediately at the temperature listed below.   |
| Stability&Storage        | Store at $\leq$ -70°C, stable for 6 months after receipt. Store at $\leq$ -70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles. |
| Reconstitution           |  |

## **SDS-PAGE** image



## Background

| Alternative Names | Cystathionine Gamma-Lyase; Cysteine-Protein Sulfhydrase; Gamma-Cystathionase;<br>CTH  |
|-------------------|---|
| Background        | Cystathionine Gamma-Lyase (CTH) belongs to the trans-sulfuration enzymes family. CTH exists as a homotetramer and interacts with CALM in a calcium- |

## Product Name: Recombinant Human CTH Catalog #: PEH0489



dependent manner. CTH breaks down cystathionine into cysteine, ammonia and 2oxobutanoate. CTH catalyzes the last step in the trans-sulfuration pathway from methionine to cysteine and has broad substrate specificity. Defects in CTH will lead to cystathioninuria, which is an autosomal recessive phenotype characterized by abnormal accumulation of plasma cystathionine.

**Note** For Research Use Only , Not for Diagnostic Use.