


Product Name: Recombinant Human IL-1b
Catalog #: PEH0886



Summary

Name	IL-1 beta/Interleukin-1 Beta/IL-1b/IL-1F2
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<0.01 EU/μg as determined by LAL test.
Construction	Recombinant Human Interleukin-1 Beta is produced by our E.coli expression system and the target gene encoding Ala117-Ser269 is expressed.
Accession #	P01584
Host	E.coli
Species	Human
Predicted Molecular Mass	17.5 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM PB , 150 mM NaCl , 3% Trehalose, 0.02% Tween 80, 4% Mannitol, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human IL-1b
Catalog #: PEH0886

Alternative Names

Interleukin-1 beta; Catabolin; IL1F2; IL1B

Background

IL1B belongs to the IL-1 family. Interleukin 1 (IL-1) is a family of polypeptide cytokines consisting of two agonists, IL-1 alpha (IL-1F1) and IL-1 beta (IL-1F2) encoded by two distinct genes and perform identical biological functions. IL-1 stimulates thymocyte proliferation by inducing IL-2 release, B-cell maturation and proliferation, and fibroblast growth factor activity. IL-1 proteins are involved in the inflammatory response. It is identified as endogenous pyrogens, and is reported to stimulate the release of prostaglandin and collagenase from synovial cells.

Note

For Research Use Only , Not for Diagnostic Use.