


Product Name: Recombinant Human IL-37
Catalog #: PEH1006



Summary

Name	IL-37/Interleukin-37
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Interleukin-37 is produced by our E.coli expression system and the target gene encoding Lys53-Asp218 is expressed.
Accession #	Q9NZH6
Host	E.coli
Species	Human
Predicted Molecular Mass	18.7 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, 2mM DTT, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Alternative Names Interleukin-37; FIL1 Zeta; IL-1X; Interleukin-1 Family Member 7; IL-1F7; Interleukin-

Product Name: Recombinant Human IL-37
Catalog #: PEH1006

Background

1 Homolog 4; IL-1H; IL-1H4; Interleukin-1 Zeta; IL-1 Zeta; Interleukin-1-Related Protein; IL-1RP1; Interleukin-23; IL-37; IL37; FIL1Z; IL1F7; IL1H4; IL1RP1
Human Interleukin family 1 Member 7 (IL1F7) is a member of the Interleukin 1 cytokine family. Five alternatively spliced transcript variants encoding distinct isoforms have been reported with distinct expression profiles. The longest IL1F7 transcript, referred to as IL1F7b or IL1F7 isoform 1, encodes a 218 amino acid residues proprotein containing a 45 amino acid propeptide, which is cleaved to generate mature protein. IL1F7b binds to IL18 R α with low affinity but does not exert any IL18 agonistic or antagonistic effects. IL1F7b also binds interleukin 18 binding protein (IL-18BP), an inhibitory binding protein of interleukin 18 (IL-18), and subsequently forms a complex with IL18 receptor beta subunit, and through which it inhibits the activity of IL-18.

Note

For Research Use Only , Not for Diagnostic Use.