
Product Name: Recombinant Human SNAP- alpha (N-6His) Catalog #: PEH1190



Summary

Name	NAPA/Alpha-soluble NSF attachment protein
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/µg as determined by LAL test.
Construction	Recombinant Human Alpha-Soluble NSF Attachment Protein is produced by our E.coli expression system and the target gene encoding Met1-Arg295 is expressed with a 6His tag at the N-terminus.
Accession #	P54920
Host	E.coli
Species	Human
Predicted Molecular Mass	35.4 KDa
Formulation	Lyophilized from a 0.2 μ m filtered solution of 20mM Tris-HCl, 150mM NaCl, pH 8.0.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Product Name: Recombinant Human SNAP- alpha (N-6His) Catalog #: PEH1190

Background

Alternative Names	Alpha-Soluble NSF Attachment Protein; SNAP-Alpha; N-Ethylmaleimide-Sensitive
	Factor Attachment Protein Alpha; NAPA; SNAPA
Background	α -Soluble NSF Attachment Protein (SNAP- α) is a member of the SNAP (Soluble
	NSF Attachment Protein) family. SNAP- α interacts with PRKCABP and disrupts the
	interaction between GRIA2 and PRKCABP, leading to the internalization of GRIA2.
	SNAP- α is required for vesicular transport between the endoplasmic reticulum and
	the Golgi apparatus. SNAP- α is in charge of the binding of NSF and therefore the
	formation of a 20S fusion particle.

Note

For Research Use Only, Not for Diagnostic Use.