Product Name: Recombinant Human NHP2L1 (N-6His) Catalog #: PEH1227



Summary

Name NHP2-like protein 1/NHP2L1

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/µg as determined by LAL test.

Construction Recombinant Human NHP2-Like Protein 1 is produced by our E.coli

expression system and the target gene encoding Met1-Val128 is expressed

with a 6His tag at the N-terminus.

Accession # P55769

Host E.coli

Species Human

Predicted Molecular Mass 16.3 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, 600mM NaCl, pH

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 Stability&Storage


months under sterile conditions after opening. Please minimize freeze-thaw

cvcles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

> not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Product Name: Recombinant Human NHP2L1 (N-6His)



Catalog #: PEH1227

Background

Alternative Names

Background

NHP2-Like Protein 1; High Mobility Group-Like Nuclear Protein 2 Homolog 1; OTK27; SNU13 Homolog; hSNU13; U4/U6.U5 tri-snRNP 15.5 kDa Protein; NHP2L1 NHP2-Like Protein 1 (NHP2L1) is a member of the ribosomal protein L7Ae family. NHP2L1 proteinis limited to the nucleus, primarily focused in the dense fibrillar component of the nucleolus. NHP2L1 has been shown to interact with RAD17and PRPF31. The protein undergoes a conformational change upon RNA-binding. NHP2L1 binds to the 5-stem-loop of U4 snRNA and may play a role in the late stage of spliceosome assembly, prior to step I of splicing catalysis.

Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838