Product Name: Recombinant Human NIP7 (N-6His)

Catalog #: PEH1228



Summary

Name NIP7/KD93

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Human 60S Ribosome Subunit Biogenesis Protein NIP7

Homolog is produced by our E.coli expression system and the target gene

encoding Met1-Thr180 is expressed with a 6His tag at the N-terminus.

Accession # Q9Y221

Host E.coli

Species Human

Predicted Molecular Mass 22.6 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, 100mM NaCl, pH

8.0.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human NIP7 (N-6His) Catalog #: PEH1228

C EnkiLife

Alternative Names 60S Ribosome Subunit Biogenesis Protein NIP7 Homolog; KD93; NIP7

Background 60S Ribosome Subunit Biogenesis Protein NIP7 Homolog (NIP7) belongs to the

NIP7 family. NIP7 contains one PUA domain, it is essential for the process of proper 27S pre-rRNA and 60S ribosome subunit assembly. NIP7 is a monomer form and interacts with NOL8 and SBDS, and may bind to RNA. In addition, NIP7 is one of the many trans-acting factors required for eukaryotic ribosome biogenesis, which interacts with nascent pre-ribosomal particles and dissociates as they

complete maturation and are exported to the cytoplasm.

Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838