


## Summary

<b>Name</b>	Phosphoglucomutase 2/PGM2
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE
<b>Endotoxin level</b>	<1 EU/μg as determined by LAL test.
<b>Construction</b>	Recombinant Human Phosphoglucomutase-2 is produced by our E.coli expression system and the target gene encoding Met1-Asp612 is expressed with a 6His tag at the N-terminus.
<b>Accession #</b>	AAH10087.1
<b>Host</b>	E.coli
<b>Species</b>	Human
<b>Predicted Molecular Mass</b>	70.5 KDa
<b>Formulation</b>	Lyophilized from a 0.2 μm filtered solution of 20mM Tris-HCl, 200mM NaCl, 0.06% Tween80, pH8.0
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
<b>Stability&amp;Storage</b>	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
<b>Reconstitution</b>	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## SDS-PAGE image



## Background

**Product Name: Recombinant Human PGM2 (N-6His)**  
**Catalog #: PEH1323**



---

**Alternative Names**

Phosphoglucomutase-2; PGM 2; Glucose phosphomutase 2; Phosphodeoxyribomutase; Phosphopentomutase

**Background**

Phosphoglucomutase-2 (PGM2) is a member of PGM family, which catalyzes the inter-conversion of sugar phosphates and participates in anabolic and catabolic reactions. When cells are grown in glucose, PGM catalyzes the conversion of glucose-6-phosphate to glucose-1-phosphate an important precursor required for the synthesis of UDP glucose and trehalose. PGM2 catalyzes the conversion of the nucleoside breakdown products ribose-1-phosphate and deoxyribose-1-phosphate to the corresponding 5-phosphopentoses, and it may also catalyze the interconversion of glucose-1-phosphate and glucose-6-phosphate. But this protein has low glucose 1,6-bisphosphate synthase activity.

**Note**

For Research Use Only , Not for Diagnostic Use.