Product Name: Recombinant Human S100A13

Catalog #: PEH1390



Summary

Name Protein S100-A13/S100A13

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Human Protein S100-A13 is produced by our E.coli expression

system and the target gene encoding Ala2-Lys98 is expressed.

Accession # Q99584

Host E.coli

Species Human

Predicted Molecular Mass 11.3 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of 50mM Tris-HCl, 1mM CaCl2, 0.1%

Tween-20, pH8.0.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human S100A13

Catalog #: PEH1390



Alternative Names Protein S100-A13;S100A13;S100 calcium-binding protein A13;

Background S100A13 is a member of the S100 family of proteins containing 2 EF-hand calcium-

binding motifs. It is widely expressed in various types of tissues with a high expression level in thyroid gland. In smooth muscle cells, this protein co-expresses with other family members in the nucleus and in stress fibers, suggesting diverse functions in signal transduction. It plays a role in the export of proteins that lack a signal peptide and are secreted by an alternative pathway. It binds two calcium ions per subunit and one copper ion. Binding of one copper ion does not interfere with calcium binding. It is required for the copper-dependent stress-induced export of IL1A and FGF1. The calcium-free protein binds to lipid vesicles containing

phosphatidylserine, but not to vesicles containing phosphatidylcholine.

Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838