## **Product Name: Recombinant Human RPS19**

Catalog #: PEH1446



#### **Summary**

Name RPS19/40S ribosomal protein S19

**Purity** Greater than 95% as determined by reducing SDS-PAGE

**Endotoxin level** <1 EU/μg as determined by LAL test.

Construction Recombinant Human 40S Ribosomal Protein S19 is produced by our E.coli

expression system and the target gene encoding Pro2-His145 is expressed.

Accession # P39019

**Host** E.coli

**Species** Human

Predicted Molecular Mass 16.1 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, 1mM EDTA, pH 7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

**Stability&Storage** Store at  $\leq$ -70°C, stable for 6 months after receipt. Store at  $\leq$ -70°C, stable for 3


months under sterile conditions after opening. Please minimize freeze-thaw

cycles

**Reconstitution** Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

### **SDS-PAGE** image



## **Background**

# **Product Name: Recombinant Human RPS19**

Catalog #: PEH1446



Alternative Names 40S Ribosomal Protein S19; RPS19

Background 40S Ribosomal Protein S19 (RPS19) is a ribosomal protein that Belongs to the

ribosomal protein S19e family. RPS19 is located in the nucleoli, and higher level expression is seen in colon carcinoma tissue than normal colon tissue. It required for pre-rRNA processing and maturation of 40S ribosomal subunits. RPS19 plays a role in many biological processes, such as endocrine pancreas development, erythrocyte differentiation, mRNA metabolic process. Defects in RPS19 are the cause of Diamond-Blackfan anemia type 1 (DBA1), which is a form of Diamond-Blackfan anemia, a congenital non-regenerative hypoplastic anemia that usually presents early in infancy. Diamond-Blackfan anemia is characterized by a moderate to severe macrocytic anemia, erythroblastopenia, and an increased risk of

malignancy.

#### Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838