


Product Name: Recombinant Mouse C3a
Catalog #: PEM0433

Summary

| | |
|---------------------------------|--|
| Name | Complement Component C3a/C3a |
| Purity | Greater than 95% as determined by reducing SDS-PAGE |
| Endotoxin level | <1 EU/ μ g as determined by LAL test. |
| Construction | Recombinant Mouse Complement Component C3a is produced by our E.coli expression system and the target gene encoding Ser671-Arg748 is expressed. |
| Accession # | P01027 |
| Host | E.coli |
| Species | Mouse |
| Predicted Molecular Mass | 9.2 kDa |
| Formulation | Lyophilized from a 0.2 μ m filtered solution of PBS, pH 7.4. |
| Shipping | The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below. |
| Stability&Storage | Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles. |
| Reconstitution | Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. |

SDS-PAGE image



Background

Product Name: Recombinant Mouse C3a
Catalog #: PEM0433

Alternative Names

Complement Component C3a; Anaphylatoxin; C3a

Background

Complement is defined as key part of innate immunity and as the first line of defense in the fight against invading pathogens. Complement 3 (C3) is the most abundant component of the complement cascade and the convergent point for all three major complement activation pathways: namely classical, alternative and mannose-binding lectin pathways. Complement activation leads to the formation of the C3 convertase, which cleaves C3 into the key effector molecules, C3a (anaphylatoxin) and C3b (opsonin) which then drive microbe removal. By binding to C3a receptor (C3aR), C3a exhibits potent anaphylatoxin activity, including increased vascular permeability, triggering degranulation of mast cells, inflammation, and activating leukocytes.

Note

For Research Use Only , Not for Diagnostic Use.