


Summary

Name	IL-2/Interleukin-2/T cell growth factor/TCGF
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<0.01 EU/μg as determined by LAL test.
Construction	Recombinant Mouse Interleukin-2 is produced by our E.coli expression system and the target gene encoding Ala21-Gln169 is expressed.
Accession #	P04351
Host	E.coli
Species	Mouse
Predicted Molecular Mass	17.4 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM Sodium Citrate, 0.2% Tween 80, pH 3.0.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Mouse IL-2
Catalog #: PEM0898

Alternative Names

aldesleukin; interleukin 2; interleukin-2; IL-2; IL2; T-cell growth factor; T cell growth factor; TCGF

Background

Interleukin 2 (IL 2), also termed T-cell growth factor, is a member of the cytokine family which includes IL-4, IL-7, IL-9, IL-15 and IL-21. Each member of this family has a four alpha helix bundle. IL-2 signals through the IL-2 receptor, a complex consisting of three subunits, termed alpha, beta and gamma. The IL-2 R gamma is shared by cytokine receptors of all members of cytokine family. Mature mouse IL/xad2 shares 56% and 73% aa sequence identity with human and rat IL/xad2, respectively. IL-2 is produced by CD4+ T cell, CD8+ T cells, gamma δ T cells, B cells, dendritic cells and eosinophils, and plays a vital role in key function of the immune system, tolerance and immunity, primarily via its potent stimulatory activity for T cells. Thus, IL/xad2 may be a key cytokine in the natural suppression of autoimmunity.

Note

For Research Use Only , Not for Diagnostic Use.