


Summary

Name	LIF/Leukemia inhibitory factor
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/ μ g as determined by LAL test.
Construction	Recombinant Mouse Leukemia Inhibitory Factor is produced by our E.coli expression system and the target gene encoding Ser24-Phe203 is expressed.
Accession #	P09056
Host	E.coli
Species	Mouse
Predicted Molecular Mass	19.9 KDa
Formulation	Lyophilized from a 0.2 μ m filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at $\leq -20^{\circ}\text{C}$, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8 $^{\circ}\text{C}$ for 2-7 days. Aliquots of reconstituted samples are stable at $\leq -20^{\circ}\text{C}$ for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Mouse LIF
Catalog #: PEM1084

Alternative Names Leukemia inhibitory factor;Differentiation-stimulating factor;lif;D factor

Background Mouse Leukemia inhibitory factor (lif) is a secreted protein which belongs to the LIF/OSM family.LIF has been implicated in a many physiological processes including development, hematopoiesis, bone metabolism, and inflammation. it has the capacity to induce terminal differentiation in leukemic cells. Its activities include the induction of hematopoietic differentiation in normal and myeloid leukemia cells, the induction of neuronal cell differentiation, and the stimulation of acute-phase protein synthesis in hepatocytes.

Note

For Research Use Only , Not for Diagnostic Use.