


Product Name: Recombinant Human Azurocidin (C-6His)
Catalog #: PHH0114



Summary

Name	HBP/Azurocidin/CAP37
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Cationic Antimicrobial Protein-37 is produced by our Mammalian expression system and the target gene encoding Ile27-Pro250 is expressed with a 6His tag at the C-terminus.
Accession #	P20160
Host	Human Cells
Species	Human
Predicted Molecular Mass	25.24 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM HEPES, 150mM NaCl, pH 7.5.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human Azurocidin (C-6His)
Catalog #: PHH0114



Alternative Names

Azurocidin; Cationic Antimicrobial Protein CAP37; Heparin-Binding Protein; HBP; AZU1

Background

Azurocidin is an Azurophil granule antibiotic protein, with monocyte chemotactic and antibacterial activity. The Azurophil granules, specialized lysosomes of the neutrophil, contain at least 10 proteins implicated in the killing of microorganisms. Azurocidin is a member of the serine protease family that includes Cathepsin G, Neutrophil Elastase (NE), and Proteinase 3 (PR3), however, Azurocidin is not a serine proteinase since the active site serine and histidine residues are replaced. Human Azurocidin together with NE and PR3 are expressed coordinately and are packaged together into azurophil granules during neutrophil differentiation. Azurocidin has been identified as a modulator of endothelial permeability and an important multifunctional inflammatory mediator. Neutrophils arriving first at sites of inflammation release Azurocidin which acts in a paracrine fashion on endothelial cells causing the development of intercellular gaps and allowing leukocyte extravasation. Azurocidin thus be regarded as a reasonable therapeutic target for a variety of inflammatory disease conditions.

Note

For Research Use Only , Not for Diagnostic Use.