Product Name: Recombinant Human BCMA (C-Fc)

Catalog #: PHH0140



Summary

Name BCMA/TNFRSF17/CD269/B cell maturation antigen

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Human B-cell Maturation Protein is produced by our

Mammalian expression system and the target gene encoding Met1-Ala54 is

expressed with a human IgG1 Fc tag at the C-terminus.

Accession # Q02223

Host Human Cells

Species Human

Predicted Molecular Mass 32.8 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Alternative Names Tumor necrosis factor receptor superfamily member 17;TNFRSF17;B-cell

Product Name: Recombinant Human BCMA (C-Fc) Catalog #: PHH0140



Background

maturation protein;CD269

B cell maturation protein (BCMA) is a type III membrane protein which belongs to the TNF receptor superfamily (TNFRSF17). BCMA contains one extracellular cysteine rich domain, as well as TACI. Human BCMA is a 184 amino acid (aa) protein consisting of a 54 aa extracellular domain, a 23 aa transmembrane domain, and a 107 aa intracellular domain. Mouse and human BCMA share 62% amino acid identity. BCMA is mainly expressed in immune organs and mature B cell lines. BCMA appears to be localized to the Golgi compartment. BCMA may play an important role in B cell development, function and regulation.

Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838