Product Name: Recombinant Human BMPR2 (C-Fc-6His) Enkilife Catalog #: PHH0162

Summary

Name BMPR-II/PPH1/BMPR2/BMP type II receptor

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Human Bone Morphogenetic Protein Receptor Type IIA is

produced by our Mammalian expression system and the target gene encoding Ser27-Ile151 is expressed with a human IgG1 Fc, 6His tag at the C-

terminus.

Accession # Q13873

Host Human Cells

Species Human

Predicted Molecular Mass 41.9 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, 5%

Trehalose, 0.06% Tween 80, pH 7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Alternative Names Bone Morphogenetic Protein Receptor Type-2; BMP Type-2 Receptor; BMPR-2;

Bone Morphogenetic Protein Receptor Type II; BMP Type II Receptor; BMPR-II;

BMPR2; PPH1

Background

Bone Morphogenetic Protein Receptor II (BMPR-II) is a Type II Serine/Threonine
Kinase that mediates cellular responses to BMPs. BMPR-II is characterized by

lacking of a GS domain, and presence of a C-terminal extension typical of type II receptors. BMPRII binds BMP2, BMP4 and BMP7 weakly in the absence of type I receptor, and the binding can be facilitated by the presence of the type I receptor, including BMPR-IA/Brk1, BMPR-IB, and ActR-I. BMPR-II plays a key role in cell growth. Defects in BMPR-II have been linked to primary pulmonary hypertension. Human and mouse BMPR-II are highly conserved and share 97 % amino acid

sequence identity.

Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838