Product Name: Recombinant Human LILRA2 (C-6His)

Catalog #: PHH0377



Summary

Name LILRA2/ILT1/CD85h

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Human Leukocyte Immunoglobulin-Like Receptor Subfamily A

Member 2 is produced by our Mammalian expression system and the target

gene encoding Gly24-Ser420 is expressed with a 6His tag at the C-terminus.

Accession # Q8N149-2

Host Human Cells

Species Human

Predicted Molecular Mass 44.8 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human LILRA2 (C-6His) Catalog #: PHH0377



Alternative Names

Background

Leukocyte Immunoglobulin-Like Receptor Subfamily A Member 2; CD85 Antigen-Like Family Member H; Immunoglobulin-Like Transcript 1; ILT-1; Leukocyte

Immunoglobulin-Like Receptor 7; LIR-7; CD85h; LILRA2; ILT1; LIR7

Leukocyte Immunoglobulin-Like Receptor Subfamily A Member 2 (LILRA2) is a single-pass type I membrane protein. LILRA2 is expressed predominantly on monocytes and B cells, and at lower levels on dendritic cells and natural killer cells. LILRA2 contains four Iq-like C2-type domains, with short cytoplasmic domains lacking an immunoreceptor tyrosine-based inhibitory motif (ITIM) and with transmembrane regions containing a charged arginine residue, may initiate

stimulatory cascades. LILRA2 does not bind class I MHC antigens.

Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838