


Summary

Name	CDH16/Cadherin-16/Kidney-specific cadherin
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Cadherin-16 is produced by our Mammalian expression system and the target gene encoding Pro18-Ala786 is expressed with a 6His tag at the C-terminus.
Accession #	O75309
Host	Human Cells
Species	Human
Predicted Molecular Mass	84.4 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human CDH16 (C-6His)
Catalog #: PHH0387



Alternative Names

CDH16; Cadherin-16; Kidney-specific cadherin; Ksp-cadherin

Background

Cadherin-16(CDH16) is a single-pass type I membrane protein which contains six cadherin domains. Mature cadherin proteins consist of a large N-terminal extracellular domain, a single membrane-spanning domain, and a small highly conserved C-terminal cytoplasmic domain. Cadherins are calcium-dependent cell adhesion proteins and may contribute to the sorting of heterogeneous cell types. They preferentially interact with themselves in a homophilic manner in connecting cells. Three calcium ions are usually bound at the interface of each cadherin domain and rigidify the connections, imparting a strong curvature to the full-length ectodomain. CDH16 is exclusively expressed in kidney, where the protein functions as the principal mediator of homotypic cellular recognition. It plays a role in the morphogenic direction of tissue development. CDH16 is composed of an extracellular domain containing 6 cadherin domains, a transmembrane region and a truncated cytoplasmic domain. However, it lacks the prosequence and tripeptide HAV adhesion recognition sequence typical of most classical cadherins.

Note

For Research Use Only , Not for Diagnostic Use.