Product Name: Recombinant Human CDO (C-6His)

Catalog #: PHH0390



Summary

Name CDO

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Human Cell Adhesion Molecule-related/down-regulated by

Oncogenes is produced by our Mammalian expression system and the target

gene encoding Asp26-Pro943 is expressed with a 6His tag at the C-terminus.

Accession # Q4KMG0

Host Human Cells

Species Human

Predicted Molecular Mass 100.4 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Product Name: Recombinant Human CDO (C-6His)

Catalog #: PHH0390



Background

Alternative Names Cell adhesion molecule-related/down-regulated by oncogenes; CDON; CDO

Background CDO (CAMrelated/down-regulated by oncogenes) is a member of the

Immunoglubulin (Ig) superfamily, Ig/Fibronectin (FN) type III repeat family of cell surface proteins. Human CDO is a type I transmembrane (TM) glycoprotein. It is synthesized as a 1287 amino acid (aa) precursor that contains a 25 aa signal sequence, a 938 aa extracellular domain (ECD), a 21 aa TM segment and a 303 aa cytoplasmic region. The ECD contains five C2-type Iglike domains, followed by three FN type III repeats. The ECD of human CDO is 85% aa identical to mouse CDO ECD. CDO is found on muscle precursor and neural progenitor cells of the embryo. It likely promotes muscle differentiation, and contributes to axon

guidance and neuronal patterning.

Note

For Research Use Only , Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838