


Summary

Name	Chitinase-3-like protein 2/CHI3L2
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Chitinase 3-Like Protein 2 is produced by our Mammalian expression system and the target gene encoding Tyr27-Leu390 is expressed with a 6His tag at the C-terminus.
Accession #	AAH11460.1
Host	Human Cells
Species	Human
Predicted Molecular Mass	41.94 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, 5% Trehalose, 5% Mannitol, 0.01% Tween 80, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Product Name: Recombinant Human CHI3L2 (C-6His)
Catalog #: PHH0403



Background

Alternative Names

Chitinase-3-Like Protein 2; Chondrocyte Protein 39; YKL-39; CHI3L2

Background

Chitinase 3-Like Protein 2 (CHI3L2) is a 39 kDa secreted glycoprotein which belongs to the glycosyl hydrolase 18 family and the most closely related to human cartilage glycoprotein 39, which promotes the growth of human synovial cells as well as skin and fetal lung fibroblasts. Highest expression of CHI3L2 is in chondrocytes, followed by synoviocytes, lung and heart. It is not detected in spleen, pancreas, and liver. CHI3L2 may also be expressed in developing brain and placenta. Increased levels of CHI3L2 have been demonstrated in synovial fluids of patients with rheumatoid or osteoarthritis as well as in some other pathologies and in malignant tumors, particularly in glioblastomas. CHI3L2 may bind glycan structure with high affinity, but not heparin. It has no chitotriosidase activity, but is likely to bind some type of glycan.

Note

For Research Use Only , Not for Diagnostic Use.