


Product Name: Recombinant Human CXCL1 (C-6His)
Catalog #: PHH0465



Summary

Name	CXCL1/C-X-C motif chemokine 1/KC/GRO alpha/CINC1/MGSA-alpha
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<0.01 EU/μg as determined by LAL test.
Construction	Recombinant Human C-X-C Motif Chemokine 1 is produced by our Mammalian expression system and the target gene encoding Ala35-Asn107 is expressed with a 6His tag at the C-terminus.
Accession #	P09341
Host	Human Cells
Species	Human
Predicted Molecular Mass	8.9 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, 5% Trehalose, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human CXCL1 (C-6His)
Catalog #: PHH0465



Alternative Names

Growth-Regulated Alpha Protein; C-X-C Motif Chemokine 1; GRO-Alpha(1-73); Melanoma Growth Stimulatory Activity; MGSA; Neutrophil-Activating Protein 3; NAP-3; CXCL1; GRO; GRO1; GROA; MGSA; SCYB1

Background

Chemokine (C-X-C motif) Ligand 1 Protein (CXCL1) is a growth factor for melanoma cells and a chemotaxin for neutrophils and a member of the CXC chemokine family that is a potent neutrophil attractant and activator and is also active toward basophils. CXCL1 is expressed by macrophages, neutrophils and epithelial cells; it has neutrophil chemoattractant activity. CXCL1 plays a critical nonredundant role in the development of experimental Lyme arthritis and carditis via CXCR2-mediated recruitment of neutrophils into the site of infection and may also have important pro-nociceptive effects via its direct actions on sensory neurons, and may induce long-term changes that involve protein synthesis.

Note

For Research Use Only , Not for Diagnostic Use.