Product Name: Recombinant Human IL-2RA (C-6His)

Catalog #: PHH0916



Summary

Name IL-2 R alpha/CD25/IL2-RA/IL-2R α/IL-2 receptor subunit alpha

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Human Interleukin-2 Receptor Subunit Alpha is produced by

our Mammalian expression system and the target gene encoding Glu22-

Cys213 is expressed with a 6His tag at the C-terminus.

Accession # P01589

Host Human Cells

Species Human

Predicted Molecular Mass 22.8 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of 10mM HEPES, 50mM NaCl, pH7.2.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than $100\mu g/ml$. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than $100\mu g/ml$. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human IL-2RA (C-6His) Catalog #: PHH0916



Alternative Names

Interleukin-2 receptor subunit alpha;CD25;p55;TAC antigen;IL2-RA;IL-2R subunit

alpha;IL-2-RA;IL-2 receptor subunit alpha

Background

Interleukin-2 receptor subunit alpha (IL2RA) is a single-pass type I membrane protein, contains 2 Sushi (CCP/SCR) domains. The interleukin 2 (IL2) receptor alpha (IL2RA) and beta (IL2RB) chains, together with the common gamma chain (IL2RG), constitute the high-affinity IL2 receptor. Homodimeric alpha chains (IL2RA) result in low-affinity receptor, while homodimeric beta (IL2RB) chains produce a mediumaffinity receptor. Normally an integral-membrane protein, soluble IL2RA has been isolated and determined to result from extracellular proteolyisis.

Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838