


Summary

| | |
|---------------------------------|--|
| Name | IL-31RB/OSMRB |
| Purity | Greater than 95% as determined by reducing SDS-PAGE |
| Endotoxin level | <1 EU/μg as determined by LAL test. |
| Construction | Recombinant Human Oncostatin-M-Specific Receptor Subunit Beta is produced by our Mammalian expression system and the target gene encoding Glu28-Ser739 is expressed with a 6His tag at the C-terminus. |
| Accession # | Q99650 |
| Host | Human Cells |
| Species | Human |
| Predicted Molecular Mass | 82.03 KDa |
| Formulation | Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.2. |
| Shipping | The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below. |
| Stability&Storage | Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles. |
| Reconstitution | Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. |

SDS-PAGE image



Background

Product Name: Recombinant Human OSMRB (C-6His)
Catalog #: PHH0922



Alternative Names

Oncostatin-M-Specific Receptor Subunit Beta; Interleukin-31 Receptor Subunit Beta; IL-31 Receptor Subunit Beta; IL-31R Subunit Beta; IL-31R-Beta; IL-31RB; OSMR; OSMRB

Background

Oncostatin-M-Specific Receptor Subunit β (OSMR β) is a 150 - 180 kDa member of the IL-6 receptor family. OSMR β associates with gp130 to form the type II OSM receptor, the receptor is responsive to OSM. Gp130 subunit is shared by other IL-6 family cytokine receptors, and OSMR β associates with gp130-like receptor (GPL) to form a receptor complex responsive to IL-31. The human OSMR β cDNA encodes a 979 amino acid (aa) precursor, the precursor includes a 27 aa signal sequence, a 712 aa extracellular domain (ECD), a 22 aa transmembrane segment, and a 218 aa cytoplasmic domain. The ECD contains one partial and one complete hematopoietin domain, an Ig-like domain, and three Fibronectin type-III domains.

Note

For Research Use Only , Not for Diagnostic Use.